



Treatment options for scab control

This information is provided by SCOPS as a guide. SCOPS does everything within its power to ensure the information is up-to-date and correct but product choices remain the responsibility of the prescriber and user. Always read the manufacturer's instructions before use. Withdrawal periods are subject to change and it is the user's responsibility to ensure withdrawal periods are adhered to.

Plunge dips for scab control	Company	Chemical	Use and protection against infestation	Meat withdrawal period
Osmonds Gold Fleece Dip	Bimeda	Diazinon	Treatment and up to four weeks protection.	49 days
Paracide 62	Animax Ltd	Diazinon	Treatment and up to four weeks protection.	70 days

Injectable products for scab control	Company	Chemical	Use and protection against infestation	Meat withdrawal period
Cydectin 1%	Zoetis	Moxidectin	Prevention - one injection protects against infestation for 28 days. Treatment - two injections 10 days apart.	70 days
Cydectin 20mg/ml LA injection (2%)	Zoetis	Moxidectin	Prevention against infection or reinfection - 60 days after one injection. Treatment - one injection in base of the ear.	104 days
Dectomax 10mg/ml Solution for Injection for Cattle and Sheep	Elanco AH	Doramectin	Treatment - one injection. Avoid contact with untreated or infected and treated flocks for 14 days.	70 days
Ecomectin 10mg/ml Solution for Injection	Eco Animal Health	Ivermectin	Treatment - two injections seven days apart.	42 days
Ivomec Classic	Merial AH	Ivermectin	Treatment - two injections seven days apart.	37 days
Noromectin 1% Multi Injection	Norbrook Labs	Ivermectin	Treatment - two injections seven days apart.	42 days
Panomec Injection for Cattle, Sheep and Pigs	Merial AH	Ivermectin	Treatment - two injections seven days apart.	37 days



Treatment options for scab control

This information is provided by SCOPS as a guide. SCOPS does everything within its power to ensure the information is up-to-date and correct but product choices remain the responsibility of the prescriber and user. Always read the manufacturer's instructions before use. Withdrawal periods are subject to change and it is the user's responsibility to ensure withdrawal periods are adhered to.

Paramectin Multi Injection	Norbrook Labs	Ivermectin	Treatment - two injections seven days apart.	42 days
Premadex 1% Injection	Downland	Ivermectin	Treatment - two injections seven days apart.	43 days
Qualimec Injection 10mg/ml Solution for Injection	Elanco AH	Ivermectin	Treatment - two injections seven days apart.	42 days
Zermex 1% Injection	Downland	Moxidectin	Prevention - one injection protects against infestation for 28 days. Treatment - two injections 10 days apart.	70 days
Zermex 20mg/ml LA for injection (2%)	Downland	Moxidectin	One injection for treatment and prevention. Persistency of reinfection - 60 days.	104 days