Example 3

Low scab risk / high liver fluke risk



www.scops.org.uk







SUNDAY	SATURDAY	FRIDAY	THURSDAY	WEDNESDAY	TUESDAY	MONDAY	
Turn out to Far Meadow where sheep have grazed until recently but now isolated from the rest of	6	Treated for S Roundworms (Group 5- SI drench (bronze) and closantel drenches given.	150 shearling ewes 4 bought direct from farms from area where liver fluke is significant issue.	3	1 2	1 2	1
the flock.	urs	used on arrival for 48 ho	Hou				
14	13	12	11	10	9	8	
21	PEC test shows no worm eggs so treatment was effective.	Take mob FEC sample to test the wormers have worked effectively and blood for the ELISA test for sheep scab.	18	17	16	15	
28	27	Carry out A Fluke FEC or Copro-antigen test here to check the efficacy of the initial treatment.	Injected with Doramectin and moved to fields not grazed by sheep for the last 15 days.	Results of scab test are positive. Treatment required.	23	22	
4	3	2	1	31	30	29	
11	10	9	Shearlings can now join the flock.	7	6	5	
18	17	Second dose of 16 closantel for liver fluke given.	15	14	13	12	

Should be given regardless of FEC/CoAg result

Example 3

Low scab risk / high liver fluke risk



Assess risk
- Roundworms

- Sheep scab

- Liver fluke

1

Treat

Yard or house animals for 24 – 48 hours 3

Isolate

Turn out on to pastures grazed previously by resident flock 3-4 weeks 4

Test

Ensure treatment was effective Is scab a possible risk?

5

Rejoin floc

Join the rest of the flock

26

Retreat

Re-treat for liver fluke if necessary www.scops.org.uk

SCOPS







MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY	SUNDAY
1	2	3	150 shearling ewes 4 bought direct from farms from area where liver fluke is significant issue.	Treated for 5 Roundworms (Group 5- SI drench (bronze) and closantel drenches given.	6	Turn out to Far Meadow 7 where sheep have grazed until recently but now isolated from the rest of the flock.
0	0	40				3
8	9	10	11	12	13	14
15	16	17	18	Take mob FEC sample 19 to test the wormers have worked effectively and blood for the ELISA test for sheep scab.	20 FEC test shows no worm eggs so treatment was effective.	21
22	23	Results of scab test are positive. Treatment required.	Injected with 25 Doramectin and moved to fields not grazed by sheep for the last 15 days.	Carry out A Fluke 26 FEC or Copro-antigen test here to check the efficacy of the initial treatment.	27	28
29	30	31	1	2	3	4
5	6	7	Shearlings can now join the flock.	9	10	11
12	13	14	15	Second dose of closantel for liver fluke given.	17	18